

ON THE GALACTIC DISK METALLICITY DISTRIBUTION FROM OPEN CLUSTERS

I. NEW CATALOGUES AND ABUNDANCE GRADIENT

L. Chen, J.L. Hou AND J.J. Wang

Shanghai Astronomical Observatory, CAS, Shanghai 200030, China

chenli@center.shao.ac.cn, hjlyx@center.shao.ac.cn

ABSTRACT

In this paper we have compiled two new open cluster catalogues. In the first one, there are 119 objects with ages, distances and metallicities available, while in the second one, 144 objects have both absolute proper motion and radial velocity data, of which 45 clusters also with metallicity data available.

Taking advantages of the large number of objects included in our sample, we present an iron radial gradient of about -0.063 ± 0.008 dex kpc $^{-1}$ from the first sample, which is quite consistent with the most recent determination of oxygen gradient by nebulae and young stars, which is about -0.07 dex kpc $^{-1}$. By dividing clusters into age groups, we show that iron gradient was steeper in the past, which is consistent with the recent result from Galactic planetary nebulae data, and also consistent with the inside-out galactic disk formation scenarios. Based on the cluster sample, we also discussed the metallicity distribution, cluster kinematics and space distribution. A disk age-metallicity relation could be implied from those properties, although we could not give conclusive result from the age metallicity diagram based on the current sample. More observations are needed for metal poor clusters. From the second catalogue, we have calculated the velocity components in cylindrical coordinates with respect to the GSR for 144 open clusters. The velocity dispersion of the older clusters are larger than that of young clusters, but they are all much smaller than that of the Galactic thick disk stars.

Subject headings: open clusters: metallicity and kinematics — Galaxy: formation — Galaxy: evolution

1. INTRODUCTION

Since the seminal work of Eggen, Lynden-Bell and Sandage (1962), great progress has been made in understanding the formation and evolution of the Milky Way galaxy. The progress comes, on one side, from observations concerning chemical abundances in stars (and clusters), gas clouds and, on the other side, from improved knowledge relevant to galaxy formation and evolution.

However, some important quantities related to the chemical evolution of our Galaxy, such as star formation history, initial mass function, gas flow, etc., are not yet well understood. Observational data from the Milky Way disk and halo has put strong constraints on our understanding of those quantities. Among a variety of observables, radial abundance gradients along the Galactic disk is one of the most important constraints on the Galactic chemical evolution model. The existence of such gradients is now well established, through radio and optical observations of HII regions, disk stars, planetary nebulae (Henry and Worthey 1999; Hou et al. 2000; Chiappini et al. 2001; Maciel et al. 2002) and open clusters (Friel 1995, 1999). An average gradient of ~ -0.06 dex kpc $^{-1}$ is observed in the Milky Way disk for most of the elements, e.g. O, S, Ne, Ar and Fe. This magnitude of the observed gradients constrains the various parameters in the chemical evolution model, such as the timescales of star formation and infall (Prantzos and Aubert 1995) or any variations of the stellar initial mass function properties with metallicities (Chiappini et al. 2001).

In the last decade, a number of successful models have been developed related to the chemical evolution of the Milky Way galaxy, but some important differences exist. One of them concerns the history of abundance gradient along the Galactic disk: were they steeper or flatter in the past? Different predictions were made by various models, although most of them claim that they could reproduce the majority of the observational properties both in the solar neighborhood and on the whole disk. Time flattening evolution is suggested by the models of Prantzos and Aubert (1995); Mollà et al. (1997); Allen et al. (1998); Boissier and Prantzos (1999); Chang et al. (1999); Hou et al. (2000); Chang et al. (2002), while the opposite is supported by models of Tosi (1988); Samland et al. (1997) and Chiappini et al. (1997, 2001).

The situation is neither settled observationally. Estimated ages of various types (PNI, PNII, PNIII) of planetary nebulae(PN) span a large fraction of the age of the Galaxy. Observations of the abundances of those objects across the Milky Way disk could, in principle, provide some information on the time evolution of the abundance gradient (Maciel and Köppen 1994; Maciel and Quireza 1999; Maciel and Costa 2002; Maciel et al. 2002). In a recent work, Hou et al. (2000) have made a detailed analysis for the O, Ne, S and Ar gradient based on the PN data of Maciel and Quireza (1999). It was shown that there is fairly good

agreement between model predictions and observations concerning all the properties of the observed abundance profiles (absolute values, gradient, scatter) for O, S, Ne and Ar. The model suggests that abundance gradients are steeper in the earlier epoch. However, the large scatter in the adopted data does not allow one to conclude on the temporal variation of the gradients. Nevertheless PNs suffer from large uncertainties concerning their progenitor’s masses and lifetimes as well as their distances from Galactic center.

On the other hand, open clusters(OCs) have long been used to trace the structure and evolution of the Galactic disk (Friel 1995). Since open clusters could be relatively accurately dated and we can see them to large distance, their $[\text{Fe}/\text{H}]$ values serve an excellent tracer to the abundance gradient along the Galactic disk as well as many other important disk properties, such as Age-Metallicity Relation(AMR), abundance gradient evolution, disk age and so on (Carraro et al. 1998).

At this point, one might ask whether the field disk populations are also able to trace the disk evolution. Indeed, the extensive studies by Edvardsson et al. (1993), and recently by Chen et al. (2000), who concentrate on disk F, G stars, show an overall radial gradient that is nearly independent of age. Those results are based on stars mainly restricted in the solar neighborhood. A more detailed analysis for the disk iron gradient was given by Cui et al. (2000) on the basis of 1302 field star with high resolution proper motion and parallax data from Hipparcos satellite. They have derived an radial iron gradient of $-0.057 \text{ dex kpc}^{-1}$ within galactocentric distance from 8.5 kpc to 17 kpc. However, it is still difficult to reveal any pronounced gradient evolution from those results. Moreover, results from those studies are strongly affected by selection effects and rely on the techniques for determining individual stellar distances (which are heavily dependent on the adopted Galaxy potential model) that are much less reliable than those used to obtain cluster distances. In a recent work, Corder and Twarog (2001) have modelled the effects of the orbital diffusion of stars and clusters on the Galactic abundance gradient. The general conclusion is that the effect of diffusion makes a gradient shallower over time, and the cluster population offers a more viable means for finding detailed structure within the recent Galactic abundance gradient.

Here, we also pointed out that our recent treatment on deriving the abundance gradient from open clusters in Hou et al. (2002) is in fact not proper. In that paper, we have simply taken four Catalogs from literatures (Carraro et al. 1998; Twarog Ashman and Anthony-Twarog 1997; Piatti et al. 1995; Friel 1995), and merge them just by making cross checking for the common clusters, without examining individually to see if there are important difference among clusters in the different catalogs (Twarog 2002). We refer this paper to act as a substitution to our previous one.

In this paper, we compiled a set of new open cluster catalogues. The catalogue was

divided into two parts: CAT 1 and CAT 2. In CAT 1, we list 119 clusters with iron abundance, age, distance and reddening data available. This could provide statistically more significant information to the Galactic disk formation and evolution, such as Age-Metallicity Relation, abundance gradient and its time or/and spatial evolution and so on. The second sample consists of 144 clusters with three dimensional kinematics information available. From this sample, we are able to explore some statistical relations among kinematics and other observables.

The paper is organized as the following: firstly in Section 2, we describe the main characteristic of the two samples. Then, in Section 3, we give some statistical analysis for the sample, mainly some metallicity and kinematics distributions. The abundance gradient is given in Section 4. We show that, based on our open cluster data, the abundance gradient of the Galactic disk was steeper in the past. In Section 5, we make some detailed discussions about the Age-Metallicity Relation (AMR) of the disk. Finally, a brief summary is given in Section 6.

2. THE CATALOGUE

During the past decades, a number of authors have presented their statistical studies on the Galactic disk based on their own open cluster catalogue. However, most of the catalogues suffer from either lacking of homogeneity in the cluster age and metallicity or insufficient three dimensional kinematic data.

With the full release of Hipparcos Catalogue (ESA 1997) and the latest Tycho 2 catalogue (Høg et al. 2000), we have seen a large growth of proper motion data for open clusters (e.g. Baumgardt et al. (2000); Dias et al. (2001)). A most recent compilation was given by Dias et al. (2002). In their catalogue, information of 1537 open clusters were present, from which 9% have both mean proper motion and radial velocity data simultaneously; 37% have distance, $E(B - V)$ and age determinations, including 96 clusters also have iron abundances data available.

We have compiled two new catalogues of the Galactic open clusters. The first one (hereafter CAT 1) lists 119 (including Berk29) clusters parameters for distance, age and metallicity. The age, distance and reddening information are all (except NGC1348, NGC2158 and Tombaugh 2) from Dias et al. (2002), while most iron abundance data (96 clusters) were taken from Dias et al. (2002). The metallicities of another 23 clusters are from other 10 literatures (Cameron 1985; Kubiak et al. 1992; Friel and Janes 1993; Friel 1995; Edvardsson et al. 1995; Piatti et al. 1995; Brown et al. 1996; Gratton 2000; Ann et al. 2002; Carraro et

al. 2002).

Thus far CAT 1 provides a most complete open cluster sample concerning the iron abundance, distance and age parameters together. This sample could provide statistically more significant information concerning the Galactic AMR, radial iron gradient as well as its evolution, etc.

In the second catalogue (hereafter CAT 2), we have listed observed kinematical data from literature for 144 clusters, *with both radial velocity and mean proper motion available*. The mean radial velocity data are mostly (122 of 144 objects) from a compilation in WEBDA database (<http://obswww.unige.ch/webda/meanvr.html>), primarily based on the work of Rastorguev et al. (1999). The absolute proper motion of 125 clusters, based on the Hipparcos system, are from Baumgardt et al. (2000). Mean proper motions of additional 16 cluster were added from compilation of Dias et al. (2001), with cluster membership probability derived by Tycho2 proper motions. Data of NGC2355 comes from Soubiran et al. (2000), and data of Coma Ber and Pleiade clusters are from Robichon et al. (1999). In fact, the above observed kinematic information constitutes a sub-catalogue of that of Dias et al. (2002). But, here in CAT 2 we have further calculated the three dimensional velocity of open clusters by combing with radial velocity and mean absolute proper motion data and give, for each cluster, the velocity components(Π, Θ, W) in cylindrical coordinates with respect to the Galactic Standard Reference (GSR). In addition, for each cluster, age and iron abundance data are also listed whenever available. (Notice that in the spatial velocity calculation, following parameters are adopted for the Sun: galactocentric distance 8.5 kpc, velocity components relative to LSR (10.0, 15.0, 8.0 km s^{-1}) and the rotation velocity : 225.0 km s^{-1}).

Our catalogue files (Table 1, Table 2) are self-explanatory. CAT 1 (Tab.1) consists of data for 119 (118+Berk29) open clusters. For each cluster, we list its heliocentric galactic coordinates in B1950.0 and the following parameters, when available: galactocentric distance; distance from the Sun; color excess $E(B-V)$, age in Gyr; mean metallicity, and two reference codes separated by a coma, where the first number is for age, distance and $E(B-V)$ data while the second number for iron abundance value. In CAT 2 (tab.2), we provide information of spatial motion for 144 clusters. For each object, the following data are listed: galactic coordinates in B1950.0; mean radial velocity ; mean proper motions; velocity components in cylindrical coordinates with respect to the GSR; mean spatial velocity and error; age and iron abundance when available.

In order to check if the data in our CAT 1 has any significant systematic difference with other published catalogues, we have made a comparison with Friel’s catalogue (Friel 1995), with 41 clusters in common. We found that the average difference in metallicity is less than 0.10dex, well within the typical observational uncertainty. The average difference

for R_{GC} is about 0.5kpc. Note that the age indicator in Friel’s work is based on the MAI (Morphological Age Index), which was only intended to provide a relative age ranking of clusters, therefore, it is not fully comparable. But there still has good overall correlation between Friel’s catalogue and ours.

Our following analysis will be mainly based on those two catalogues, but excluding cluster Berkeley 29. In CAT 1, Berkeley 29 has the galactocentric radius of 23 kpc, the $E(B - V)$ of 0.15 and a metallicity of -0.18 dex, from the compilation of Dias et al. (2002). However, different values for these parameters of Berkeley 29 were published in the literature. Kaluzny (1994) gave a much smaller galactocentric distance of about 19 kpc, the reddening $E(B - V)$ larger than 0.21 and based on the CMD morphology and comparison to other old clusters, he also deduced a $[Fe/H]$ value of lower than -1. In the work of Noriega-Mendoza and Ruelas-Mayorgo (1997), who applied a new technique for simultaneous determination of $[Fe/H]$ and $E(B - V)$, a $[Fe/H]=-0.30$ and a $E(B - V) = 0.01$ were given for Berkeley 29. As the properties of Berkeley 29 are quite uncertain we do not include this object in the following calculations. In general, the uncertainty for the metallicity determinations in open clusters is about 0.1dex.

3. STATISTICAL PROPERTIES

3.1. The Galactic distribution of open clusters

Using data from Dias et al. (2002) for 571 open clusters with distance and age data, we plotted the cluster positions on an (X,Y) coordinate system, with the zero point in X at the galactic center (the Sun is assumed to be at 8.5 kpc) as Figure 1 shows. Here the full line arc represents the solar circle about the galactic center. One will find from this figure that in the galactic plane, young clusters (with ages younger than that of Hyades, 0.8 Gyr, see Phelps et al. (1994)) distributed quite uniformly around the Sun, while roughly only 20% of the old clusters are inside the solar circle, most of the old ones are located further away from the galactic center than the Sun. This result is quite consistent with the early comprehensive study of Phelps et al. (1994). The deficiency of older clusters in the inner part of the disk has been ascribed to the preferential destruction of the clusters when they encountered with giant molecular clouds, which were primarily found in the inner Galaxy.

The distribution of either old or young clusters perpendicular to the galactic plane could be fitted by a simple exponential law, which are plotted in Figure 2. The younger clusters are distributed on the galactic plane almost symmetrically about the Sun, with a perpendicular scale height of approximately 57 pc. In contrast, about 80% old clusters are in the outer

disk, outside $R_{gc}=10$ kpc; this population has a scale height of about 354 pc. The derived scale heights are in excellent agreement with the early results of Janes et al. (1988), Janes and Phelps (1994). Janes and Phelps (1994) divided the open clusters into young and old components according to cluster’s MAI, and derived their scale heights as 55pc and 375pc, respectively. The remarkable agreement shows that the MAI could really be a good age indicator for open clusters.

The main advantage of our CAT 2 is that we have both radial velocity and mean proper motion available for 144 clusters. This provides a chance to probe the velocity projection on the Galactic plane, as shown in Figure 3. Obviously, most of the clusters are located in the solar neighborhood and have the velocity vectors well follow the Galactic rotational pattern.

3.2. The metallicity distribution

The metallicity distribution of 118 open clusters is plotted in the upper panel of Figure 4. Here the iron abundance of about 3/4 of the *OC* sample (with $[Fe/H] > -0.2$) has roughly a Gaussian distribution which peaks at the solar value. Meanwhile a metal poor metallicity tail is also clearly seen. Here we divide our *OC* sample by two groups, that is, Metal-Poor (hereafter MP) component and Metal-Rich (hereafter MR) component, with the dividing line between them, somewhat arbitrarily, at $[Fe/H] = -0.2$. In the lower panel of Figure 4, we show two histograms for the open clusters of the above two groups. By assuming an exponential law, we can derive their scale-heights to be 535 pc and 106 pc, for the MP and MR components, respectively. Taking the scale heights of the Galactic thick and thin disk as 760 pc and 260 pc (Ojha et al. 1996), we could see that spatially, the MP group might be either within the tail of the thick disk or in the outskirts of the thin disk while the MR group is just a thin disk component. However, our *OC* metallicity sample is surely not a complete one, and is subject to a variety of observation effects. For example, the outer disk clusters are subject to significant selection effects - they can be seen more readily if they are at larger distances from the plane, and they are likely to be older, since younger clusters which live closer to the plane will not be as visible, and more difficult to observe.

This can also be seen clearly in Fig.5, where we have plotted the dependence of clusters vertical height on the galactocentric distance for the whole sample with distances data available (Fig.5a) and for those also have metallicity data (Fig.5b). In Fig. 5b, we see that most of the MP *OCs* are outer disk objects (with a median $R_{gc} \sim 11.3$ kpc and the majority(65%) are older than 0.8 Gyr, see also Fig. 8,9) and relatively far away from the Galactic plane, with a median distance about 326 pc. For MR clusters, most of them are in the inner disk ($R_{gc} \leq 10$ kpc), distributing in the immediate solar neighborhood (with a median $R_{gc} \sim 8.7$

kpc), and about 78% are young (with ages < 0.8 Gyr) objects (also see Fig. 8,9). This phenomena is very likely to imply the possible existence of the age metallicity relation in open clusters.

Meanwhile, when $R_{gc} < 10$ kpc, most *OCs*, no matter MP or MR, are much closer to the Galactic plane, with a median height $z \sim 84$ pc. Especially, there is few clusters observed in the region of $R_{GC} < 6$ kpc. This has been attributed to the destructive power of the large numbers of giant molecular clouds in the inner regions of the Galaxy (van der Bergh and MMcClure 1980). There is much evidence that leads us to believe that open clusters have been selectively destroyed near the plane of the disk and only those clusters whose orbits keep them away from the Galactic plane can survive long enough to appear as outer disk, or metal-poor clusters. In the meanwhile, this also imply that part of the outer clusters (they are metal-poor and with high- z) might be formed as a result of disturbances to the Galactic disk, possibly caused by tidal interactions with other galaxies or infalling gas, as suggested by Janes and Phelps (1994). However, it is still hard to understand why there are almost no high z clusters in the region of $R_{GC} \sim 6.5$ -8 kpc, compared with the outer disk results.

As a comparison, both globular and open cluster metallicity distribution are plotted in the upper and lower panels of Figure 6. We can see a clear overlap between metal rich globular cluster and metal poor open clusters around $[\text{Fe}/\text{H}] \sim -0.4$ dex. If the age metallicity relation do exist, then this could be another evidence which support the idea that a possible connection between the halo and disk population exist, both in their chemical and dynamical history.

Based on a survey of proper motion stars, Carney et al. (1990) pointed out that the Galactic halo population had the chemical and dynamical history almost independent of the disk. However, from a study of the oldest open clusters, Phelps et al. (1994) found that the oldest open clusters (Be17 with an age about 12.5Gyr) have the ages compatible with that of the youngest globular clusters, suggesting that there may have been little or no delay in time between formation of the halo and the onset of the development of the disk. Phelps's argument rests largely on the age of Be17 clusters, however, recent works have given an age of about 9 Gyr to this clusters (Carraro et al. 1999), and so there does still appear to be a gap between the formation of the halo and the thick disk. On the other hand, if the cluster metallicity is related to their age, the metallicity overlapping of the clusters, as we have presented, may be another indication of the connection proposed by Phelps et al. (1994). In fact, the distinction between "open" and "globular" (so-called "super") clusters may turn out to be largely an artificial one (Larsen 2002). They could be both originated from the Super Star Clusters (SSCs) which are observed in large numbers in interacting galaxies and merger remnants. Our Milky Way disk is very likely undergone a process of minor mergers

in the early epoch. The thick disk is plausibly the result of heating of the thin disk through such events (Wyse 2001).

3.3. Metallicity vs. kinematic

In Figure 7, we plot the relation between cluster rotational velocities around the Galactic center, Θ , and the galactocentric distance. There exhibits, from least-square fitting, an insignificant slope of about $-2.5 \text{ km s}^{-1} \text{ kpc}^{-1}$, with quite large scatter. In the right panel, we present the dependence of velocity dispersion on the cluster age. The young clusters have a smaller velocity dispersion, which is expected from their small z scale height. Although the velocity dispersion for the older clusters is about 20 km s^{-1} , it is still much smaller than that of the Galactic thick disk stars, which is about 50 km s^{-1} . Anyway, for most of the CAT.2 clusters (90%), their heights from the galactic plane are well within 200pc, they are just thin disk objects.

In our kinematic sample, there are only 2 objects in the outer disk (Berkeley 31 and Dolidze 25, both with $R_{gc} > 14 \text{ kpc}$). They have unreliable proper motion and radial velocity results (with relative errors up to about 50%) and thus were not included in the above radial gradient fitting.

4. DISK METALLICITY GRADIENT FROM OPEN CLUSTERS

4.1. The abundance gradients

The first radial metallicity gradient using open clusters was given by Janes (1979) based on DDO and UBV photometric data of 41 disk objects (part of them are field stars). The derived gradient is $-0.05 \text{ dex kpc}^{-1}$. Panagia and Tosi (1981), by matching theoretical isochrones to HR diagrams of 20 clusters with age less than 1 Gyr, derived an iron abundance gradient of $-0.095 \text{ dex kpc}^{-1}$. A similar result was also obtained by Cameron (1985) based on 37 clusters with mixed ages. By introducing a weighting system in order to evaluate and compare the published parameters in Lyngå (1987) Catalog of Open Clusters data, Janes et al. (1988) determined some basic parameters of 413 open clusters, such as ages, distances, linear diameters and so on. Among them, 87 open clusters have the metallicity data. They have derived a gradient about $-0.133 \text{ dex kpc}^{-1}$. By separating the clusters into age groups, they found that young clusters have much smaller gradient than that of older clusters. Besides, all those authors found some indications that the gradient became shallower in the direction of the Galactic center and steeper in the outer parts of the Galaxy.

Friel and Janes (1993), (hereafter FJ93) presented their results from a spectroscopic study of a sample of giant stars in 24 open clusters. They derived a galactocentric radial abundance gradient of $[\text{Fe}/\text{H}]$ about $-0.088 \text{ dex kpc}^{-1}$. A subsequent revision of the FJ93 result was presented by Friel (1995), using additional spectroscopic results and a more uniform set of cluster properties. From a sample of 44 clusters, Friel (1995) derived an iron gradient of $-0.091 \text{ dex kpc}^{-1}$. At the same time, Piatti et al. (1995) derived a much smaller gradients, $-0.07 \text{ dex kpc}^{-1}$, from a sample of 63 open clusters with a wide range of ages. These results are quite consistent with the recent result of Friel (1999) who obtained a gradient about $-0.06 \text{ dex kpc}^{-1}$. Another gradient result was presented by Carraro et al. (1998) recently. The metallicities of all selected 37 clusters were obtained spectroscopically. The final gradient was about $-0.085 \text{ dex kpc}^{-1}$, agreed with earlier result of FJ93. By dividing the sample into age bins, it was found that the present-day gradient is a little shallower than the past one, while the middle epoch seems to display a steepening of the gradient.

The presence of a linear gradient for open clusters has been questioned by Twarog Ashman and Anthony-Twarog (1997) (hereafter TAA97). TAA97 put forth an alternative description, namely, step function, about the radial abundances distribution of the open clusters. Within this work, a set of 76 clusters with abundances based upon DDO and/or moderate dispersion spectroscopy has been transformed to a common metallicity scale and used to study the local structure and evolution of the Galactic disk. They found that the metallicity distribution of clusters with galactocentric distance is best described by two distinct zones, with a sharp discontinuity at $R_{GC} = 10 \text{ kpc}$. Between $R_{GC} = 6.5 \text{ kpc}$ and 10 kpc , the clusters have a mean metallicity of 0.0 dex with, at best, weak evidence for a shallow gradient over this range, while those beyond 10 kpc have a mean value about -0.30 dex . This two-step distribution seems quite similar with the nebula results of Simpson et al. (1995). Neglecting this two-step phenomena, a least square fitting results in a gradient about $-0.067 \text{ dex kpc}^{-1}$ between 6 and 15 kpc if cluster BE21 was excluded because both metallicity and distance of this object are quite uncertain.

The existence of radial iron abundance gradients is also confirmed by our new up-to-date sample. The result is shown in the upper panel of Figure 8. By equal-weighted least-square fitting, we derived a radial abundance gradient of $-0.063 \pm 0.008 \text{ dex kpc}^{-1}$, which agrees well with most of the previous open cluster results. And it is also similar to the gradients obtained from other tracers, such as disk HII regions and planetary nebulae (see a summary in Hou et al. (2000)). The existence of gradient along the galactic disk provides good opportunity to test theories of disk evolution and stellar nucleosynthesis. It suggests that the role of the Galactic bar in inducing large scale radial mixing and therefore flattening the gradient has been rather limited; alternatively, the bar could be too young ($< 1 \text{ Gyr}$) to have brought any important modifications to the gaseous and abundance profile. However, we

must notice that our current knowledge on the iron gradient as derived from open clusters is far from being clear. Open clusters span a wide range of age, from several millions years to several Gyr, therefore they do not trace the young component of the galactic disk. The result we obtained is somewhat an averaged one (over age). The obtained similarity of gradient between iron and other elements, such as oxygen, is quite surprising since the sites of nucleosynthesis for iron and oxygen are quite different. It is well known that iron is mainly produced in type SNIa, while oxygen is largely a product of SNII, that is from massive stars. So the abundance history is very different for those two types of elements. The gradient similarity might be simply coincidental, or further investigations should be given in the production nature of those elements.

We also derived a vertical abundance gradient of -0.295 ± 0.050 dex kpc⁻¹ (lower panel of Figure 8). This is consistent with the result of Carraro et al. (1998).

4.2. Gradient evolution in the Galactic disk

As we have pointed out in Section 1, the behavior of gradient evolution along the Galactic disk is a major problem for different chemical evolution models. Open cluster system is an ideal template for this analysis because *OCs* have relatively well determined ages, distances and metallicities.

In the upper panel of Figure 9, we show gradients for two sub-samples with cluster age < 0.8 Gyr (80 clusters) and ≥ 0.8 Gyr (38 clusters), respectively. The fitting results are -0.024 ± 0.012 dex kpc⁻¹ for younger clusters, -0.075 ± 0.013 dex kpc⁻¹ for older ones. If we take the mean age for the youngest and oldest clusters as 0.00 Gyr and 6.00 Gyr (this is somewhat arbitrary, just for illustration purpose) in our sample, we can estimate an average flattening rate of 0.008 dex kpc⁻¹Gyr⁻¹ during the past 6 Gyr. Similar value is obtained by Maciel et al. (2002) from PN data for [O/H].

As we have indicated in the Introduction, that the time evolution of the abundance gradient along the Galactic disk is crucial in discriminating different theoretical models that adopt various prescriptions used for the time dependence of the SFR and the infall. Our current open clusters sample could surely provide some insights on this subject. The time flattening tendency we obtained supports the 'inside-out' disk formation scenarios with infall time scale dependent on radius from the disk center (Boissier and Prantzos 1999; Hou et al. 2000; Chang et al. 2002).

In the lower panel of Figure 9, we divided clusters into inner (< 10 kpc) and outer groups. The corresponding gradients are -0.040 ± 0.022 dex kpc⁻¹ and -0.047 ± 0.023

dex kpc^{-1} , respectively. We can see that the inner disk exhibits roughly the same (or a bit smaller) gradient as the outer part. This result is also consistent with the abundance gradient determined by using Cepheid in the solar neighborhood (Andrievsky et al. 2002). However, in our CAT 1, the inner most cluster is located at a galactocentric distance about 6.8 kpc, it is necessary to have more inner clusters data (between 3 kpc and 7 kpc) in order to further check the gradient behavior for the inner disk. If the Galactic bar does play the role, then the inner gradient could be more flat compared with outer part.

Our cluster sample is nearly 50 % more than that of TAA97, and we did not find evidence of any abrupt discontinuity. A similar conclusion was reached by Friel (1999), using high resolution abundance determinations for metallicity calibration.

5. DISK AGE-METALLICITY RELATION

The age-metallicity relation (AMR) for the Galactic disk provides useful clues about the chemical evolution history of the Milky Way, and also put an important constraint on the theoretical models of the disk. The observed abundance data generally show a decrease of the stellar metallicity with increasing stellar age, indicating a continuous growth of the metals in the ISM during the life of the Galaxy. The early study on AMR for nearby stars by Twarog (1980) found that the mean metallicity of the disk increased by a factor of five between 12 and 5 billion year ago and has increased only slightly since then. This was also confirmed by latter photometric survey of Meusinger et al. (1991). With the high resolution spectroscopic data, Edvardsson et al. (1993) showed a plot of iron abundance versus relative ages for the 189 stars in the solar neighborhood. The overall trend of a slowly increasing abundance with decreasing age was consistent with the previous photometric results. However, the most striking feature of their result is the large scatter around the average trend, which marks a weak correlation between age and metallicity. This spread was, as they pointed out, in part due to selection bias for the programme stars, and at least partly intrinsic, since the mean errors in $[\text{Fe}/\text{H}]$ measurement and logarithmic age derivation are much less than the scatter.

In a recent paper, Feltzing et al. (2001) have re-examined the Galactic AMR in the solar neighborhood based on a sample of 5828 dwarfs and sub-dwarfs from Hipparcos Catalogue. They found that the solar neighborhood age-metallicity diagram is well populated at all ages and especially that old, metal-rich stars do exist, which have been omitted in previous samples. This indicates a complete lack of enrichment over the age of Galactic disk among the fields stars in the solar neighborhood.

Using open clusters to explore the AMR has the main advantage both in abundance and

age determinations since one is dealing with a group of stars and the result is less susceptible to individual errors (Carraro et al. 1998). Cameron (1985) was the first to probe the AMR from open cluster data, and found no age-metallicity relation based on his cluster sample. This is not surprising since the metallicity of the Galactic disk increased only slightly during the past 5 Gyr, while his sample of 38 clusters contained no objects older than 5.1 Gyr. More recently, Carraro et al. (1998) compiled a relatively homogenous sample of 37 open clusters. The data have more expanded cluster ages up to 9 Gyr. After correcting for the radial abundance gradient, the derived AMR showed similar trend to that of nearby stars.

In this paper, we have present a new open cluster catalogue with much more objects. The results, based on this larger sample, would be statistically more reliable. As we have shown in the Sect 3.2, statistically, the space distributions (scale heights for metal poor and metal rich groups, for young and old clusters) of open clusters are very likely imply the existence of age-metallicity relation in the Galactic disk. In Fig. 10, we plot the dependence of metallicity on the cluster age, after correcting the radial metallicity gradient. Unfortunately, it is difficult to draw any conclusive indication for AMR based on this plot due to the deficiency of very older clusters. More observational efforts should be added in finding more older clusters.

The significant spread of the AMR seems real, but its origin is not yet clear. For the scatter in the AMR of nearby stars, many possible causes have been suggested, such as orbital diffusion of stars, inhomogeneous chemical enrichment in the Galaxy evolution, overlapping of different galactic substructures and so on. All the above mentioned effects may contribute the observed scatter, while for open clusters, the result should not be very sensitive to orbital diffusion effects (Corder and Twarog 2001). Therefore, the scatter of AMR along the Galactic disk from both clusters and field disk stars is an essential feature in the formation and evolution of the Milky Way.

6. Summary

The main work of this paper is to compile a most complete open clusters sample with metallicity, age, distance data as well as kinematic information available. And upon this sample, some statistical analysis on spatial and metallicity distributions have been made.

We derived an iron radial gradient about $-0.063 \pm 0.008 \text{ dex kpc}^{-1}$ from the CAT 1, which is quite consistent with the most recent determination of oxygen gradient in nebulae and young stars. By dividing clusters into age groups, we show that iron gradient was steeper in the past, which is consistent with the recent result from Galactic planetary nebulae data. Our result supports the inside-out Galactic disk formation mechanism that invoking SFR

and infall time scale various with radius.

The spatial variation of gradient was also explored. When the clusters were divided into inner and outer groups, we found that the abundance gradient was a bit shallower inside 10 kpc. But the inner most cluster in our sample is in $R_{GC} = 6.8\text{kpc}$, we need more data of clusters in the inner region. This could be helpful to judge the radial flow effect on the current galactic chemical evolution model.

From scale heights of metal poor and metal rich clusters, we noticed that the metallicity could be related to the age. However, by plotting directly the dependence of cluster abundances on their ages, no strike slope in AMR was found. However, the paucity of metallicity of very old open clusters made it impossible to give a definite conclusion based on the current sample.

The Authors thank Drs. Chenggang Shu, Ruixiang Chang and Prof. Chenqi Fu for their helpful discussion. The authors would also thanks to an anonymous referee for invaluable comments and stimulating criticism of the manuscript, which greatly improved this article. This research was supported partly by the National Natural Science Foundation of China (No.19873014, No.10173017, No.10133020, No.19833010) and NKBRSG 1999075404 & 406, and partly by SRF for ROCS, SEM.

REFERENCES

- Allen, C., Carigi, L., & Peimbert, M. 1998, ApJ, 494, 247
- Andrievsky, S. M., Kovtyukh, V. V., Luck, R. E., Lèpine, J. R.D., Bersier, D., et al. 2002, A&A, 381, 32
- Ann, H. B., Lee, S. H., Sung, H., Lee, M. G., Kim, S. -L. et al. 2002, AJ, 123, 905
- Baumgardt, H., Dettbarn, C., & Wielen, R. 2000, A&A, 146, 251
- Boissier, S., & Prantzos, N. 1999, MNRAS, 307, 857
- Brown, J. A., Wallerstein, G., Geisler, D., & Oke, J. B. 1996, AJ, 112, 1551
- Cameron, L. M. 1985, A&A, 147, 47
- Carney, B. W., Latham, D. W., & Laird, J. B. 1990, AJ, 99, 572
- Carraro, G., Ng, Y. K., & Portinari, L. 1998, MNRAS, 296, 1045

- Carraro, G., Vallenari, A., Girardi, L., & Richichi, A. 1999, *A&A*, 343, 825
- Carraro, G., Girardi, L., & Marigo, P. 2002, *MNRAS*, 332, 705
- Chang, R. X., Hou, J. L., Shu, C.G. & Fu, C. Q. 1999, *A&A*, 350, 38
- Chang, R. X., Shu, C.G., & Hou, J.L. 2002, *CJAA*, 2, 226
- Chen, Y. Q., Nissen, P. E., & Zhao, G. 2000, *A&AS*, 141, 491
- Chiappini, C., Matteucci, F., & Gratton, R. 1997, *ApJ*, 477, 765
- Chiappini, C., Matteucci, F., & Romano, D. 2001, *ApJ*, 550, 1044
- Clariá, J.J., & Mermilliod, J.-C. 1992, *AApSS*, 95, 429
- Corder, S., & Twarog, B. A. 2001, *AJ*, 122, 895
- Cui, C. Z., Zhao, G., Zhao, Y. H., & Shi, J. R. 2000, *Science in China (series A)*, 30, 953
- Dias, W. S., Lépine, J. R., & Alessi, B. S. 2001, *A&A*, 376, 441
- Dias, W. S., Alessi, B. S., Moitinho, A., & Lépine, J. R. 2002, *A&A*, 389, 871
- Edvardsson, B., Andersen, J., Gustafsson, B., et al. 1993, *A&A*, 275, 101
- Edvardsson B., Pettersson B., Kharrazi M., et al. 1995, *A&A*, 293, 75
- Eggen, O. J., Lynden-Bell, D., & Sandage, A. R. 1962, *ApJ*, 136, 748
- ESA, 1997, *The Hipparcos and Tycho Catalogues* (European Space Agency (ESA))
- Feltzing, S., Holmberg, J. & Hurley, J.R. 2001, *A&A*, 377, 911
- Friel, E. D. 1993, *PASP*, 101, 244
- Friel, E. D. 1995, *ARA&A*, 33, 381(F95)
- Friel, E. D. 1999, *Ap&SS*, 265, 271
- Friel, E. D., & Janes, K. A. 1993, *A&A*, 267, 75 (FJ93)
- Gratton, R. 2000, *ASPC*, 198, 225
- Carney, B. W., and Harris, W. E. 2001, *Star Clusters: Saas-Fee Advanced Course 28*, ed. L.Labhardt and B.Binggeli (Springer, Swiss Society for Astrophysics and Astronomy)

- Henry, R. B. C., & Worthey, G. 1999, *PASP*, 111, 919
- Høg, E., Fabricius, C., Makarov, V. V., Urban, S., Corbin, T., et al. 2000, *A&A*, 355, L27
- Hou, J. L., Prantzos, N., & Boissier, S. 2000, *A&A*, 362, 921
- Hou, J. L., Chang, R.X., & Chen, L. 2002, *CJAA*, 2,17
- Janes, K. A. 1979, *ApJS*, 39, 135
- Janes, K. A., Tilley, C., & Lyngå, G. 1988, *AJ*, 95, 771
- Janes, K. A., & Phelps, R. L. 1994, *AJ*, 108, 1773
- Kaluzny, J. 1994, *A&AS*, 108, 151
- Kubiak, M., Kaluzny, J., Krzeminski, W., & Mateo, Mario, *Acta Astronomica*, 1992, 42, 155
- Larsen, S. S. 2002, *IAU Symposium series*, 207 (in press)
- Lyngå, G. 1987, *Computer Based Catalogue of Open Cluster Data*, 5th ed. (Strasbourg: CDS)
- Maciel, W. J., & Köppen, J. 1994, *A&A*, 282, 436
- Maciel, W. J., & Quireza, C. 1999, *A&A*, 345, 629
- Maciel, W. J., & Costa, R. D. D. 2002, *IAU Symp.* 209, in press (astro-ph/0112210)
- Maciel, W. J., Costa, R. D. D., & Uchida, M. M. M. 2002, *A&A*, accepted (astro-ph/0210470)
- Mermilliod, J.-C., Andersen, J., Nordström, B., & Mayor, M. 1995, *A&A*, 299, 53
- Mermilliod, J.-C., Mayor, M., & Burki, G. 1987, *A&AS*, 70, 389
- Mermilliod, J.-C., Hustamendia, G., del Rio, G., & Mayor M. 1996, *A&A*, 307, 80
- Mermilliod, J.-C., & Mayor, M. 1989, *A&A*, 219, 125
- Mermilliod, J.-C., & Mayor, M. 1990, *A&A*, 237, 61
- Meusinger, H., Stecklum, B., & Reimann, H.-G. 1991, *A&A*, 245, 57
- Mollà, M., Ferrini, F., & Diaz, A. I. 1997, *ApJ*, 475, 519
- Noriega-Mendoza, H., & Ruelas-Mayorga, A. 1997, *AJ*, 113, 722

- Ojha, D. K., Bienayme, O., Robin, A. C., Creze, M., & Mohan, V. 1996, *A&A*, 311, 4560
- Panagia, N., & Tosi, M. 1981, *A&A*, 96, 306
- Phelps, R. L., Janes, K. A., & Montgomery, K. A. 1994, *AJ*, 107, 1079
- Piatti, A., Claria, J. J., & Abadi, M. G. 1995, *AJ*, 110, 2813
- Piatti, A. E., Clariá, J. J., Bica, E., Geisler, D., & Minniti, D. 1998, *AJ*, 116, 801
- Portinari, L., & Chiosi, C. 2000, *A&A*, 355, 929
- Prantzos, N., & Aubert, O. 1995, *A&A*, 301, 69
- Raboud, D., & Mermilliod, J.-C. 1998, *A&A*, 329, 101
- Rastorguev, A. S., Glushkova, E. V., Dambis, A. K., & Zabolotskikh, M. V. 1999, *AZh Pisma*, 25, 689 (English transl. *Astron. Lett.*, 25, 595)
- Robichon, N., Arenou, F., Mermilliod, J.-C., & Turon, C. 1999, *A&A*, 345, 471
- Samland, M., Hensler, G., & Theis, Ch. 1997, *ApJ*, 476, 544
- Scott, J. E., Friel, E. D., & Janes, K. A. 1995, *AJ*, 109, 1706
- Simpson, J. P., Colgan, S. W. J., Rubin, R. H., Erickson, E. F., & Haas, M. R. 1995, *ApJ*, 444, 721
- Soubiran, C., Odenkirchen, M., & Le Campion, J.-F. 2000, *A&A*, 357, 484
- Tosi, M. 1988, *A&A*, 197, 47
- Twarog, B. A. 1980, *ApJ*, 242, 242
- Twarog, B. A. 2002, private communication
- Twarog, B. A., Ashman, K. A., & Anthony-Twarog, B. J. 1997, *AJ*, 114, 2556 (TAA97)
- van der Bergh, S., & McClure, R.D. 1980, *A&A*, 88, 360
- Wyse, R. F. G. 2001, *ASPC*, 230, 71




Fig. 1.— Spacial distribution of the open clusters on the galactic plane. The open circles are old clusters with age great than that of Hyades (0.8 Gyr), and the dots are for younger ones. The Sun is at $X = 8.5, Y = 0$ kpc. The galactic center is at $(0,0)$. The circle has a radius of 8.5 kpc, centered on the galactic center.




Fig. 2.— Number distribution from the galactic plane, z , for old and young sub-groups of open clusters. The fitted scale heights for the two groups are 354pc and 57pc, respectively.




Fig. 3.— Velocity projection on the Galactic plane for 144 clusters, which have both radial velocity and mean proper motion available. The Sun is located at $X = 8.5$ kpc, $Y = 0$ pc.




Fig. 4.— Upper panel: metallicity distribution of the Galactic open clusters. Lower panel: number distribution from the galactic plane, Z , for metal poor and metal rich sub-groups of open clusters. We can see that the two groups are different in their scale properties, with scale height about 535 pc and 106 pc, respectively.




Fig. 5.— Upper panel: height from the galactic plane vs. galactocentric distance, for all 571 clusters with data available. Lower panel: same as the upper panel, but only for 118 *OCs* with also iron abundance data. It can be seen that the most metal poor clusters are located in the outer part of the Galactic disk.




Fig. 6.— Comparison of the metallicity distributions for Galactic globular and open clusters




Fig. 7.— Left: correlation between clusters rotational velocity Θ and galactocentric distance. Right: velocity dispersion vs. cluster age.




Fig. 8.— Radial (upper panel) and vertical (lower panel) abundance gradient for 118 open clusters. The least-square fitting results in a gradient of $-0.063 \pm 0.008 \text{ dex kpc}^{-1}$ and $-0.295 \pm 0.050 \text{ dex kpc}^{-1}$, respectively. The typical error bar for $[\text{Fe}/\text{H}]$ is about 0.1dex, as showed in lower left corner of the figures. When deriving the vertical gradient, the radial gradient has been corrected.




Fig. 9.— Upper panel: Time evolution of the iron gradient. Triangles show clusters with age less than 0.8 Gyr, stars show clusters with age greater than 0.8 Gyr. The gradients are $-0.024 \text{ dex kpc}^{-1}$ and $-0.075 \text{ dex kpc}^{-1}$, respectively. Lower pane: gradients for inner disk (within 10 kpc) and out disk clusters. The corresponding gradients are $-0.040 \text{ dex kpc}^{-1}$ and $-0.047 \text{ dex kpc}^{-1}$, respectively.




Fig. 10.— Age-Metallicity Relation (AMR) for the 118 open clusters after correcting for the radial gradient. The solid line is a least-square fitting for the open cluster data.

Table 1. Age, Distance and Metallicity of Open Clusters

No.	Name	l (deg)	b (deg)	R_{GC} (kpc)	R_{SUN} (pc)	E(B-V) mag	Age (Gyr)	[Fe/H] dex	refs
1	Berkeley 12	161.66	-1.99	11.5	3162	0.7	4	0.07	1,1
2	Berkeley 17	175.65	-3.65	11.1	2700	0.7	12.00	-0.33	1,1
3	Berkeley 18	163.63	5.01	14.1	5800	0.46	4.26	0.02	1,1
4	Berkeley 19	176.9	-3.59	13.3	4831	0.4	3.09	-0.50	1,2
5	Berkeley 20	203.51	-17.28	16.3	8400	0.12	6.02	-0.75	1,3
6	Berkeley 21	186.84	-2.51	13.4	5000	0.76	2.18	-0.83	1,1
7	Berkeley 22	199.8	-8.05	15.8	7663	0.7	1.06	-0.30	1,1
8	Berkeley 23	192.6	5.44	15.3	6918	0.4	0.79	0.07	1,1
9	Berkeley 29	197.98	8.02	23.0	14871	0.157	1.05	-0.18	1,1
10	Berkeley 31	206.25	5.12	16.3	8272	0.08	2.05	-0.50	1,1
11	Berkeley 32	207.95	4.4	11.3	3100	0.16	3.38	-0.58	1,1
12	Berkeley 39	223.46	10.09	12.3	4780	0.12	7.94	-0.17	1,1
13	Berkeley 60	118.84	-1.63	11.2	4365	0.86	0.16	0.07	1,1
14	Berkeley 64	131.91	4.6	11.5	3981	1.05	1	-0.61	1,1
15	Berkeley 70	166.89	3.58	12.5	4158	0.48	4	-0.32	1,1
16	Berkeley 104	117.63	1.22	11.2	4365	0.45	0.79	0.07	1,1
17	Blanco 1	15.57	-79.26	8.4	269	0.01	0.06	0.23	1,4
18	Collinder 140	244.97	-7.91	8.6	405	0.03	0.03	-0.10	1,1
19	Collinder 261	301.68	-5.52	7.5	2190	0.27	8.9	-0.14	1,1
20	IC 2391	270.36	-6.9	8.5	175	0.008	0.04	-0.09	1,1
21	IC 2581	284.59	0.03	8.2	2446	0.415	0.01	-0.34	1,1
22	IC 2602	289.6	-4.9	8.4	161	0.024	0.03	-0.09	1,5
23	IC 2714	292.4	-1.79	8.1	1238	0.341	0.34	-0.01	1,1
24	IC 4651	340.08	-7.9	7.6	888	0.116	1.14	0.09	1,1
25	IC 4725	13.7	-4.43	7.9	620	0.476	0.09	0.17	1,1
26	IC 4756	36.38	5.24	8.1	484	0.192	0.5	-0.06	1,1
27	King 5	143.74	-4.27	10.0	1900	...	1	-0.38	1,1
28	King 6	143.36	-0.07	9.2	871	0.5	0.25	0.46	1,1
29	King 8	176.39	3.12	14.8	6403	0.58	0.41	-0.46	1,1
30	King 11	117.16	6.47	10.1	2892	1.27	1.11	-0.23	1,1

Table 1—Continued

No.	Name	l (deg)	b (deg)	R_{GC} (kpc)	R_{SUN} (pc)	E(B-V) mag	Age (Gyr)	[Fe/H] dex	refs
31	King 15	120.74	-0.92	10.4	3162	0.7	0.25	0.07	1,1
32	Melotte 20	146.93	-7.11	8.6	185	0.09	0.07	-0.05	1,5
33	Melotte 22	166.57	-23.52	8.6	150	0.03	0.13	-0.03	1,5
34	Melotte 25	180.06	-22.34	8.5	45	0.01	0.78	0.13	1,5
35	Melotte 66	259.55	-14.24	10.1	4313	0.143	2.78	-0.35	1,1
36	Melotte 71	228.96	4.49	10.8	3154	0.113	0.23	-0.30	1,1
37	Melotte 111	221.35	84.02	8.5	96	0.013	0.44	-0.05	1,5
38	NGC 1039	143.63	-15.6	8.8	499	0.07	0.17	-0.30	1,1
39	NGC 1193	146.74	-12.19	12.2	4300	0.12	7.9	-0.29	1,1
40	NGC 1245	146.64	-8.92	10.9	2876	0.3	0.5	0.10	1,1
41	NGC 1342	154.95	-15.34	9.0	665	0.319	0.45	-0.16	1,1
42	NGC 1348	146.96	-3.7	10.0	1820	1.02	0.13	0.07	6,6
43	NGC 1545	153.36	0.18	9.1	711	0.303	0.28	-0.06	1,1
44	NGC 1662	187.71	-21.11	8.9	437	0.304	0.42	-0.09	1,1
45	NGC 1817	186.13	-13.12	10.4	1972	0.334	0.4	-0.26	1,1
46	NGC 188	122.78	22.46	9.6	2047	0.082	4.28	-0.02	1,1
47	NGC 2099	177.64	3.09	9.8	1383	0.302	0.34	0.09	1,1
48	NGC 2141	198.08	-5.81	12.3	4033	0.25	1.7	-0.26	1,1
49	NGC 2158	186.64	1.78	12.1	3600	0.55	2.0	-0.23	7,1
50	NGC 2168	186.59	2.19	9.3	816	0.262	0.09	-0.16	1,1
51	NGC 2204	226.01	-16.1	10.4	2629	0.085	0.78	-0.33	1,1
52	NGC 2243	239.5	-17.97	11.3	4458	0.051	1.07	-0.44	1,1
53	NGC 2251	203.58	0.1	9.7	1329	0.186	0.26	-0.08	1,1
54	NGC 2259	201.76	2.07	11.6	3311	0.59	0.32	0.07	1,1
55	NGC 2281	174.9	16.88	9.0	558	0.063	0.35	0.13	1,1
56	NGC 2287	231.01	-10.44	8.9	693	0.027	0.24	0.04	1,1
57	NGC 2301	212.56	0.29	9.2	872	0.028	0.16	0.06	1,1
58	NGC 2304	197.2	8.89	12.3	3991	0.1	0.79	-0.32	1,1
59	NGC 2335	223.62	-1.18	9.5	1417	0.393	0.16	-0.03	1,1
60	NGC 2343	224.32	-1.17	9.2	1056	0.118	0.01	-0.3	1,1

Table 1—Continued

No.	Name	l (deg)	b (deg)	R_{GC} (kpc)	R_{SUN} (pc)	E(B-V) mag	Age (Gyr)	[Fe/H] dex	refs
61	NGC 2355	203.3	11.8	10.5	2200	0.12	0.7	-0.07	1,1
62	NGC 2360	229.8	-1.43	9.8	1887	0.111	0.56	-0.15	1,1
63	NGC 2420	198.11	19.63	11.3	3085	0.029	1.11	-0.26	1,1
64	NGC 2423	230.48	3.54	9.0	766	0.097	0.73	0.14	1,1
65	NGC 2437	231.87	4.07	9.4	1375	0.154	0.24	0.05	1,1
66	NGC 2477	253.59	-5.83	8.9	1222	0.279	0.7	0.01	1,1
67	NGC 2482	241.63	2.03	9.2	1343	0.093	0.4	0.12	1,1
68	NGC 2489	246.71	-0.78	10.7	3957	0.374	0.01	0.08	1,1
69	NGC 2506	230.57	9.92	11.0	3460	0.081	1.1	-0.37	1,1
70	NGC 2516	273.81	-15.85	8.4	409	0.101	0.11	0.06	1,1
71	NGC 2527	246.08	1.85	8.7	601	0.038	0.44	-0.09	1,8
72	NGC 2539	233.73	11.12	9.3	1363	0.082	0.37	0.13	1,1
73	NGC 2546	254.9	-1.98	8.7	919	0.134	0.07	0.12	1,1
74	NGC 2547	264.45	-8.53	8.5	455	0.041	0.03	-0.16	1,1
75	NGC 2548	227.93	15.39	9.0	769	0.031	0.36	0.08	1,1
76	NGC 2567	249.79	2.94	9.2	1677	0.128	0.29	-0.09	1,8
77	NGC 2571	249.09	3.54	9.0	1342	0.137	0.03	0.05	1,9
78	NGC 2632	205.92	32.48	8.6	187	0.009	0.72	0.14	1,1
79	NGC 2660	265.85	-3.03	9.1	2826	0.313	1.07	-0.18	1,1
80	NGC 2682	215.66	31.91	9.1	908	0.059	2.56	0.00	1,1
81	NGC 2818	261.98	8.58	8.9	1855	0.121	0.42	-0.17	1,1
82	NGC 2972	274.73	1.75	8.5	2062	0.343	0.09	-0.07	1,1
83	NGC 3114	283.34	-3.83	8.3	911	0.069	0.12	0.02	1,1
84	NGC 3532	289.55	1.34	8.3	486	0.037	0.31	-0.02	1,1
85	NGC 3680	286.78	16.92	8.2	938	0.066	1.19	-0.09	1,1
86	NGC 381	124.93	-1.22	9.2	1148	0.4	0.32	0.07	1,1
87	NGC 3960	294.36	6.18	7.8	2258	0.302	0.66	-0.17	1,1
88	NGC 4349	299.71	0.82	7.6	2176	0.384	0.2	-0.12	1,8
89	NGC 5138	307.54	3.53	7.4	1986	0.262	0.09	0.12	1,1
90	NGC 5316	310.22	0.12	7.7	1215	0.267	0.15	-0.02	1,8

Table 1—Continued

No.	Name	l (deg)	b (deg)	R_{GC} (kpc)	R_{SUN} (pc)	E(B-V) mag	Age (Gyr)	[Fe/H] dex	refs
91	NGC 5822	321.7	3.59	7.8	917	0.15	0.66	-0.02	1,1
92	NGC 6025	324.54	-5.88	7.9	756	0.159	0.07	0.19	1,9
93	NGC 6067	329.75	-2.21	7.3	1417	0.38	0.11	0.13	1,1
94	NGC 6087	327.76	-5.41	7.7	891	0.175	0.09	-0.01	1,1
95	NGC 6134	334.91	-0.19	7.6	913	0.395	0.92	0.18	1,1
96	NGC 6208	333.75	-5.76	7.6	939	0.21	1.17	-0.03	1,8
97	NGC 6253	335.45	-6.26	7.1	1510	0.2	5	0.36	1,1
98	NGC 6259	341.98	-1.52	7.5	1031	0.498	0.21	0.02	1,1
99	NGC 6281	347.73	1.97	8.0	479	0.148	0.31	0.00	1,8
100	NGC 6405	356.59	-0.77	8.0	487	0.144	0.09	0.06	1,1
101	NGC 6425	357.94	-1.6	7.7	778	0.399	0.02	0.09	1,8
102	NGC 6475	355.85	-4.52	8.2	301	0.103	0.29	0.03	1,8
103	NGC 6494	9.89	2.83	7.8	628	0.356	0.29	0.09	1,1
104	NGC 6633	36.1	8.3	8.2	376	0.182	0.42	0.00	1,1
105	NGC 6705	27.31	-2.78	6.8	1877	0.426	0.2	0.14	1,1
106	NGC 6716	15.39	-9.59	7.7	789	0.22	0.09	-0.31	1,9
107	NGC 6791	69.96	10.91	8.5	5853	0.117	4.39	0.15	1,1
108	NGC 6819	73.97	8.48	8.1	2360	0.238	1.49	0.07	1,1
109	NGC 6939	95.9	12.3	8.7	1185	0.32	2.21	0.02	1,1
110	NGC 6940	69.9	-7.14	8.2	770	0.214	0.72	0.01	1,1
111	NGC 7082	91.19	-2.9	8.6	1442	0.237	0.17	-0.01	1,1
112	NGC 7142	105.34	9.48	9.0	1686	0.397	1.88	0.04	1,1
113	NGC 7209	95.49	-7.33	8.6	1168	0.168	0.41	-0.12	1,8
114	NGC 752	137.18	-23.35	8.8	457	0.034	1.12	-0.08	1,1
115	NGC 7789	115.48	-5.37	9.7	2337	0.217	1.71	-0.08	1,1
116	Pismis 4	262.86	-2.43	8.5	593	0.013	0.03	-0.20	1,8
117	Ruprecht 18	239.92	-4.94	9.0	1056	0.7	0.04	-0.01	1,1
118	Ruprecht 46	238.36	5.9	8.9	752	0.07	3.98	-0.04	1,1
119	Tombaugh 2	232.83	-6.88	13.3	6300	0.4	4.0	-0.36	10,11

We list 119 clusters, while in the paper, Berkley 29 was not included in our calculations.

References. — (1) Dias et al.2002; (2) Friel and Jane 1993; (3) Friel 1995; (4) Edvardsson et al. 1995; (5) Gratton 2000; (6) Ann et al. 2002; (7) Carraro, Girardi, and Marigo 2002; (8) Piatti, Claria, and Abadi 1995; (9) Cameron 1985; (10) Kubiak et al. 1992; (11) Brown et al. 1996.

Table 2. Kinematic Parameters of Open Clusters

No.	ID	l	b	V_r	ref	$\mu_\alpha \cos \delta$	μ_δ	ref	R_{GC}	R_{SUN}	Π	Θ	W	V	σ_v	age	[Fe/H]
-----	----	---	---	-------	-----	--------------------------	--------------	-----	----------	-----------	-------	----------	-----	-----	------------	-----	--------

References. — (1) Rastorguev et al.(1999); (2) Baumgardt et al.(2000); (3) Dias et al.(2002); (4) Mermilliod et al.(1996); (5) Friel(1993); (6) Lyngå(1987); (7) Soubiran (2000); (8) Mermilliod & Mayor(1990); (9) Mermilliod & Mayor(1989); (10) Mermilliod et al. (1995); (11) Mermilliod et al.(1987); (12) Clariá & Mermilliod(1992);(13) Scott et al.(1995); (14) Raboud & Mermilliod(1998); (15) Robichon et al.(1999).

Note. — This table is available only on-line as a machine-readable table.